

SGT-003 Gene Therapy Stabilizes the DAPC and Improves Muscle Integrity in Duchenne Muscular Dystrophy

J. Patrick Gonzalez¹, Jamie L. Marshall¹, Stephen Bradley¹, Michael W. Lawlor², Tatyana A. Vetter², Brandon Chan¹, David Wolfson¹, Peter Nadeau¹, Kevin Whittlesey¹, Jessie Hanrahan¹, Gabriel Brooks¹, Matthew Harmelink¹, Nicolas Christoforou¹


¹Solid Biosciences Inc., Charlestown, MA, USA; ²Diverge Translational Science Laboratory, Milwaukee, WI, USA



ABSTRACT

Introduction: Duchenne muscular dystrophy is caused by loss of functional dystrophin, which results in destabilization of the dystrophin-associated protein complex (DAPC).^{1,2} Disruption of the DAPC weakens sarcolemmal integrity, leading to membrane fragility, chronic myofiber damage, and persistent activation of degeneration-regeneration cycles.³


SGT-003 Reconstitutes the DAPC



The DAPC anchors nNOS to the sarcolemma. Loss of dystrophin results in mislocalization of nNOS to the cytosol, leading to impaired activity-dependent nitric oxide signaling and associated vasodilation. This contributes to functional ischemia and increased susceptibility to contraction-induced injury.

In Duchenne, chronic muscle injury drives repeated degeneration-regeneration cycles, leading to elevated proportions of eMHC-positive regenerating fibers relative to healthy muscle.^{5,6}

Overview of Membrane Damage and Regeneration



Restoring expression and localization of the entire protein complex, especially the sarcoglycans and nNOS, is a critical therapeutic goal for re-establishing DAPC function and preventing severe muscle disease in Duchenne.

Objective: To evaluate whether SGT-003, Solid's next-generation adeno-associated virus (AAV) microdystrophin gene therapy, restores the DAPC to the sarcolemma, improves sarcolemmal integrity, and normalizes the degeneration-regeneration process.


Methods: Sarcolemmal localization of DAPC components (sarcoglycans, dystrobrevin, nNOS) were assessed in muscle samples from mdx mice and SGT-003-101 (INSPIRE DUCHENNE) clinical trial participants collected pre- and post-SGT-003 administration. Serum creatine kinase (CK) and titin fragments were evaluated as biomarkers of sarcolemmal integrity. eMHC positivity was measured as a marker of active degeneration-regeneration.

RESULTS

SGT-003 RESTORES THE DAPC

SGT-003 Stabilizes α-, β-, γ-, δ-Sarcoglycans; Critical DAPC Components

Figure 1. Microdystrophin-mediated reassembly of the sarcoglycan complex in mdx mice¹¹



Quadriceps tissues were harvested 33 days after dosing. Jess analysis was performed to quantify sarcoglycan protein levels. Mean microdystrophin percent positive fibers are 90% in the mdx Microdystrophin-5 treated animals. WT N=3/group, mdx Vehicle N=9/group, mdx 7E13 vg/kg N=10/group. One way ANOVA with multiple comparisons performed, asterisk p<0.05. Microdystrophin-5: microdystrophin construct used in SGT-003.

REFERENCES


1. Gao QQ, McNally EM. *Compr Physiol*. 2015;5(3):1223-39. 2. Tarakci H, Berger J. *Front Biosci (Landmark Ed)*. 2016;21(4):744-756. 3. Shi W, et al. *Muscle Nerve*. 2004;29(3):409-419. 4. Sears et al. *Circ Res*. 2003;92(5): e52-9. 5. Schiaffino S, et al. *Skelet Muscle*. 2015;5:22. 6. Guiraud S, et al. *Hum Mol Genet*. 2019;28(2):307-319. 7. Dubuisson N, et al. *Int J Mol Sci*. 2022;23(24):16080. 8. Cardone N, et al. *Acta Neuropathol Commun*. 2023; 11(1):167. 9. Forcina L, et al. *Cells*. 2020;9(5):1297. 10. Abdel-Sama E, et al. *Acta Myot*. 2009;28(3):94-100. 11. Marshall J, et al. Presented at New Directions, June 7-10, Orlando, FL. Clinical Data cutoff February 23, 2026. Solid Biosciences.

ACKNOWLEDGMENTS

This study was sponsored by Solid Biosciences Inc. (Charlestown, MA, USA).

RESULTS


Figure 2. Microdystrophin restores and co-localizes with sarcoglycans and dystrobrevin in mdx mice¹¹



Quadriceps tissues were harvested 33 days after dosing. Double stain immunofluorescence analyses were performed to assess co-localization of microdystrophin and α-, β-, γ-, δ-sarcoglycan and dystrobrevin. N=2/group. Microdystrophin-5: microdystrophin construct used in SGT-003; μDys: microdystrophin; WT: wild type.

SGT-003 Clinical Data: DAPC Restoration


Figure 3. In SGT-003 Treated Participants, microdystrophin appropriately localizes to the sarcolemma where it restores the DAPC, including sarcoglycans and dystrobrevin¹¹



SGT-003 Restores Sarcolemmal nNOS Signaling – a Key Biological Function of Dystrophin

The DAPC serves as a scaffold that localizes nNOS to the sarcolemma, enabling nitric oxide-dependent signaling as a major downstream functional consequence of DAPC integrity


Figure 4. SGT-003 microdystrophin uniquely restores membrane localization of enzymatically active, functional nNOS in mdx mice¹¹



Quadriceps tissues were harvested 92 days after dosing. Immunofluorescence analysis was performed to assess microdystrophin membrane localization and an enzymatic activity assay was performed to assess functional nNOS activity at the muscle membrane. N=10/group. nNOS: neuronal nitric oxide synthase; WT: wild type; μDys: microdystrophin.

SGT-003 Clinical Data: Reconstitution of Functional nNOS at Sarcolemma

Figure 5. SGT-003 microdystrophin uniquely restores membrane localization of functional nNOS activity in SGT-003 treated participants¹¹



SGT-003 PROTECTS MUSCLE FROM INJURY

SGT-003 Restores Sarcolemmal Integrity and Protects Against Myofiber Degeneration as Evaluated by eMHC Quantification

Figure 6. Reduction of ongoing dystrophic myofiber degeneration and regeneration observed after treatment¹¹





Figure 7. Reduction of serum biomarkers of membrane instability observed after SGT-003 treatment¹¹



SGT-003 Clinical Data: Reduces Muscle Degeneration

Figure 8. SGT-003 stabilizes the muscle membrane and prevents muscle injury in treated participants¹¹





Figure 9. SGT-003 stabilizes the muscle membrane and reduces serum markers of membrane instability in treated participants¹¹



CONCLUSIONS

- SGT-003 produces compelling biological evidence of DAPC restoration, enhanced sarcolemmal resilience, and a measurable reduction in ongoing muscle injury in both treated mdx mice and patients with Duchenne
- SGT-003 restores sarcolemmal nNOS signaling, a key biological function of dystrophin
- Critically, the decrease in eMHC suggests that muscle fibers are no longer trapped in a perpetual injury-repair loop — a shift that may enable true structural preservation and improved long-term function for patients with Duchenne
- SGT-003 has been generally well tolerated in the 53 participants dosed in the INSPIRE DUCHENNE study as of August 4, 2026